



TAMPINES MERIDIAN JUNIOR COLLEGE

JC2 PRELIMINARY EXAMINATION

CANDIDATE
NAME

CIVICS GROUP

H2 CHEMISTRY

Paper 3 Free Response

9729/03

22 September 2021

2 hours

Candidates answer on Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name and Civics Group in the spaces at the top of the page.

Write in dark blue or black pen on the answer booklet.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Section A

Answer **all** questions.

Section B

Answer **one** question.

A *Data Booklet* is provided.

The use of an approved scientific calculator is expected, where appropriate.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiners' Use		
Section A	Q1	/15
	Q2	/21
	Q3	/24
Section B	Q4	/20
	Q5	/20
Total	/80	
Grade		

Section A

Answer **all** the questions from this section.

- 1 The *rate of reaction* between an unknown bromoalkane, RBr, and aqueous sodium hydroxide, NaOH, undergoing hydrolysis to form an alcohol can be investigated experimentally using phenolphthalein as an indicator.

A few drops of phenolphthalein were added to a sample of hot aqueous sodium hydroxide. An excess of bromoalkane was added to the mixture, shaken, and the relative rate was measured when the end of the reaction was observed.

The experiment was then repeated with various initial concentrations of the bromoalkane and aqueous sodium hydroxide. All experiments were carried out at the same temperature.

- (a) (i) Explain what you understand by the term '*rate of reaction*'. [1]
 (ii) Describe how the end of the experiment would be determined. [1]

.....

.....

.....

.....

.....

.....

Table 1.1 below shows the results for the series of hydrolysis experiments.

Table 1.1

experiment	[RBr] / mol dm ⁻³	[NaOH] / mol dm ⁻³	relative rate
1	0.120	0.0400	0.0238
2	0.040	0.0400	0.00794
3	0.048	0.0800	0.00952
4	0.180	0.0200	x

- (b) (i) Deduce the order of reaction for both RBr and NaOH. [2]
 (ii) Hence write the overall rate equation for the reaction. [1]
 (iii) Predict the relative rate, **x**, for the reaction in experiment 4. [1]
 (iv) It was determined that 20% of the RBr in experiment 1 was reacted in 4.6 s. Predict the time taken for the same percentage of RBr to be reacted in experiment 3, giving your reasoning. [2]





(c) (i) Given that the identity of the bromoalkane is an aromatic chiral compound with the molecular formula C_8H_9Br , suggest a possible structure for the bromoalkane. [1]

(ii) The hydrolysis of the bromoalkane proceeds via a 2-step mechanism.

Name and describe the mechanism for the hydrolysis of the bromoalkane in aqueous sodium hydroxide. Show relevant lone pairs and dipoles, and use curly arrows to indicate the movement of electron pairs. [3]

[illegible]

- (iii) Based on the stability of the carbocation formed, suggest briefly why the bromoalkane proceeds via the mechanism proposed in (c)(ii). [1]
- (iv) Upon analysis of the final reaction mixture, it was found that the reaction mixture contained several organic side-products along with the production of the alcohol. One of the side-products was an alkene, while the other contained an ether (R–O–R) functional group.

Suggest the structures of these two side-products. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[Total: 15]



- 2 *Ketamine* is a medication that is used to induce anaesthesia. It can relieve pain in humans and animals.

The following scheme shows a possible synthesis route of *ketamine* from 1-bromo-2-chlorobenzene. Study Fig. 2.1 carefully and answer the questions that follow.




Fig. 2.1

- (a) (i) Suggest the type of reaction that occurs in steps 2, 3 and 5. [3]
- (ii) Suggest the identity of reagent U in step 5. [1]

.....


.....

.....

.....

.....


.....



- (i) Draw the skeletal formula of compound **P**. [1]
- (ii) Describe a simple chemical test to distinguish compounds **N** and **P**. State clearly the observations for each compound in the test. [2]


[illegible]

- (e) Step 2 of the reaction involves the use of a Grignard reagent (organomagnesium halide) on cyclohexanone to generate the tertiary alcohol via a nucleophilic addition mechanism.




- (i) Suggest the structural formula of the final organic product formed when

1. $(\text{CH}_3)_2\text{CHMgBr}$ reacts with CH_3CHO


2.  reacts with CO_2

[2]

- (ii) It is also possible for a Grignard reagent to react with epoxides (3-membered cyclic ethers) via a substitution reaction.



Suggest an explanation for the major product formed in the reaction below.



[1]

- (iii) Draw the mechanism between cyclohexanone and HCN with cold NaOH . Show relevant lone pairs and dipoles, and use curly arrows to indicate the movement of electron pairs.

[2]



compound	pK_b
ethylamine	3.20
ammonia	4.75
<i>ketamine</i>	6.50
phenylamine	9.30

- (i) Give an explanation for the different pK_b value for ethylamine, ammonia and phenylamine. [2]
- (ii) Suggest a reason why the pK_b value for *ketamine* is so much higher than that for ethylamine. [1]

This image shows a full page of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page, providing a template for handwriting practice or general writing. There are no margins, text, or other markings on the page.

BLANK PAGE



- 3 The organisation of the Periodic Table can be used to derive relationships between various element properties. It is also used to predict chemical properties and behaviours of undiscovered elements.

(a) Fig. 3.1 shows the third ionisation energy of eight consecutive elements (**A** to **H**) in Periods 2 and 3 of the Periodic Table.




Fig. 3.1

- (i) Write an equation for the third ionisation energy of element **B**. [1]
- (ii) Using information from Fig. 3.1, identify element **E** and write its electronic configuration. [1]
- (iii) Based on the electronic configuration of the species, explain the difference in the third ionisation energies of element **F** and element **G**. [1]
- (iv) Draw and label the orbital in which the third electron is removed from in element **G**. [1]

The Periodic Table shows helium is placed at the top of Group 18.

- (v) Suggest why helium could be placed at the top of Group 2. [1]
- (vi) Suggest why helium is not placed at the top of Group 2, by comparing **one** physical property related to its structure. [1]



After filtration of the electrolyte, the deposit underneath the anode weighed 0.09 g. On adding an excess of dimethylglyoxime to the electrolyte, 0.68 g of a bright red solid complex with the formula $\text{Ni}(\text{C}_4\text{H}_7\text{N}_2\text{O}_2)_2$ was precipitated.

(i) Calculate the expected increase in mass of copper at the cathode. [2]

(ii) Calculate the actual mass of copper removed from the impure copper strip. [2]


(iii) Hence, or otherwise, determine the percentage purity by mass of the copper strip at the anode. [1]

(iv) Suggest how the experimental procedure for collection of the solids after the filtration of the electrolyte can be improved to increase the accuracy of the value calculated in (b)(iii). [2]

[illegible]



- (c) Copper is an unreactive metal that does not react directly with dilute acids. Thus, the enthalpy change of reaction, ΔH_r , as shown below, cannot be determined directly.



In order to determine a value for ΔH_r , the enthalpy change of reaction for two reactions are needed. One of these reactions involves adding a reactive metal M to dilute sulfuric acid.



The other reaction is a displacement of copper from aqueous copper(II) sulfate, $\text{CuSO}_4(\text{aq})$, using metal M.



The value for the enthalpy change of the displacement reaction ΔH_2 can be determined using a simple calorimeter shown in Fig 3.2. The following experiment was carried out and the metal M used in this experiment can be assumed to be in excess.




Fig 3.2

In the experiment, 15.9 g of metal M was added into 50.9 cm³ of 0.300 mol dm⁻³ $\text{CuSO}_4(\text{aq})$ in a polystyrene cup. The temperature of the resulting solution increased from 25.0 °C to 60.0 °C.

The polystyrene cup used in the experiment can also absorb heat, hence a separate experiment was performed and the heat capacity of the polystyrene cup was found to be 9.7 J K⁻¹.

- (i) Based on the given information, calculate the value of ΔH_2 in kJ mol⁻¹.

You may take the density of the solution to be 1.00 g cm⁻³, and the specific heat capacity of the solution to be 4.18 J g⁻¹ K⁻¹. [2]

- (ii) Hence, or otherwise, determine the value of ΔH_r , given that ΔH_1 is -466.9 kJ mol⁻¹. [2]

reaction	E° / V
$\text{Ag}^+ + \text{e}^- \rightleftharpoons \text{Ag}$	+0.80
$\text{Na}^+ + \text{e}^- \rightleftharpoons \text{Na}$	-2.71
$\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu}$	+0.34
$2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2$	+0.00
$\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$	-2.38

Based on the list above, magnesium was chosen to be the most suitable metal M in both reactions to find ΔH_1 and ΔH_2 . Give two reasons to explain why magnesium was chosen instead of silver and sodium. [2]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

The enthalpy change of some reactions are given in Table 3.4.

$\text{CO}_2(\text{g}) + \text{O}^{2-}(\text{g}) \rightarrow \text{CO}_3^{2-}(\text{g})$	$-1148 \text{ kJ mol}^{-1}$
lattice energy of $\text{NiO}(\text{s})$	$-3870 \text{ kJ mol}^{-1}$
lattice energy of $\text{NiCO}_3(\text{s})$	$-2980 \text{ kJ mol}^{-1}$

- (i) Define the term *lattice energy*. [1]
- (ii) Explain why the lattice energy of NiO is more exothermic than that of NiCO₃. [1]
- (iii) Construct a fully labelled energy cycle and hence calculate the standard enthalpy change of thermal decomposition of NiCO₃. [3]

[illegible]




BLANK PAGE



Section B


Answer **one** question from this section.

- 4 Dinitrogen tetroxide, N_2O_4 , and nitrogen dioxide, NO_2 , exist in dynamic equilibrium with each other.



- (a) (i) Explain what is meant by *dynamic equilibrium*. [1]
- (ii) 0.0500 mol of N_2O_4 was placed in a sealed vessel of volume 1.00 dm^3 and at a temperature of 50°C . The resultant equilibrium mixture had a pressure of $1.68 \times 10^5 \text{ Pa}$, and a mass of 4.600 g.

Calculate the average relative molecular mass, M_r , of the resulting equilibrium mixture. [2]
- (iii) In this equilibrium mixture, the mole fraction of NO_2 is 0.400. Determine the degree of dissociation, α , of the N_2O_4 sample at equilibrium. [2]
- (iv) Write the expression for the equilibrium constant, K_p , for this equilibrium. [1]
- (v) Calculate the equilibrium constant, K_p at 50°C . [2]
- (vi) 0.0500 mol of N_2O_4 was placed in in a larger vessel at the same temperature, and allowed to reach equilibrium. Suggest, with reasoning, whether there will be any change to the equilibrium composition compared to the original system. [1]
- (vii) The value of pV/RT is plotted against p for one mole of an ideal gas and one mole of NO_2 gas at 50°C , where p is the pressure and V is the volume of the gas.



On the same axes, sketch the variation of pV/RT against p for one mole of NO_2 at 100°C . Briefly explain your answer. [2]





- (b) Compound **S** can be synthesised from compound **N**, (2-hydroxyphenyl)ethanal, by a four-step synthesis route shown in Fig. 4.1.




Fig. 4.1

- (i) State the reagents and conditions required for each step and suggest structures for the organic compounds **P**, **Q** and **R**. [7]
- (ii) A chemist carried out step 1 of the synthesis. After leaving the reactants to react for several hours, he wanted to determine whether the reaction had reached completion.

Suggest a simple chemical test that could be carried out for this purpose. State the expected observation if the reaction is complete. [2]



- 5 Three separate samples of white powder containing MgO , Al_2O_3 and SiO_2 were incorrectly labelled.

- (a) (i) State the pH of the resultant solution when MgO is dissolved in water. [1]
- (ii) Describe **two** simple chemical tests that could be used to distinguish the three oxides from one another. State the expected observation for each oxide in each test. [3]

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

- (b) Magnesium hydroxide, $\text{Mg}(\text{OH})_2$, is sparingly soluble in water and is used in suspension as either an antacid or a laxative.




- (i) Write an expression for the K_{sp} of $\text{Mg}(\text{OH})_2$ stating its units. [1]
- (ii) The pH of a saturated aqueous solution of $\text{Mg}(\text{OH})_2$ at $25\text{ }^{\circ}\text{C}$ is 10.35. Calculate the solubility of $\text{Mg}(\text{OH})_2$ at $25\text{ }^{\circ}\text{C}$. [2]
- (iii) Hence, or otherwise, calculate the value of K_{sp} of $\text{Mg}(\text{OH})_2$ at $25\text{ }^{\circ}\text{C}$. [1]
- (iv) State and explain how the solubility of $\text{Mg}(\text{OH})_2$ is affected by the addition of a small amount of HCl solution. [2]



	decomposition temperature / °C
Mg(OH) ₂	332
Ca(OH) ₂	512
Ba(OH) ₂	800

[illegible]



- (i) Suggest reagents and conditions for step 1. [1]
- (ii) Suggest the structures of organic products **U** and **V**. [2]
- (iii) Step 2 represents one of the possible initial steps in the synthesis of chloramphenicol. State the type of reaction in step 2. [1]



[illegible]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.



