



TEMASEK
JUNIOR COLLEGE

PRELIMINARY EXAMINATIONS

HIGHER 2

CHEMISTRY

Paper 3 Free Response

9729/03

16 September 2019

2 hours

Candidates answer on separate booklet.

Additional Materials: 12-Page Answer Booklet

Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your Civics Group, Centre number, index number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

Section A

Answer **all** questions.

Section B

Answer **one** question.

A Data Booklet is provided.

The use of an approved scientific calculator is expected, where appropriate.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **14** printed pages and **2** blank pages.

Section A

Answer **all** the questions from this section.

- 1 The term *chelates*, originates from the Greek word *chele* for “claw”. It refers to compounds containing ligands bonded to a central metal atom or ion at two or more points.

Ligand exchange occurs when ammonia and ethylenediamine (en), $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$, are added separately to copper sulfate solution.

- (a) (i) Explain all the changes observed when aqueous ammonia is added dropwise to $\text{Cu}^{2+}(\text{aq})$ till in excess. Suggest the formulae of all relevant copper compounds formed and write equations for all reactions. [4]

- (ii) Suggest why ethylenediamine can function as a *chelating ligand*. [1]


- (iii) When ethylenediamine is added to $\text{Cu}^{2+}(\text{aq})$, the following occurs.



Predict whether ethylenediamine or ammonia gives a more positive entropy change when added to $\text{Cu}^{2+}(\text{aq})$. [2]

- (iv) Suggest the coordination number and shape of $[\text{Cu}(\text{en})_2(\text{H}_2\text{O})_2]^{2+}$. [1]

- (v) The complex ion $[\text{Cu}(\text{en})_2(\text{H}_2\text{O})_2]^{2+}$ can exist as three stereoisomers, **A**, **B** and **C**. Isomers **A** and **B** rotate plane-polarised light but **C** does not. The structure of **A** is represented below.




Isomer **A**

where $\text{N} \text{---} \text{N}$ represents ethylenediamine (en)

Draw the structure of **C**.


[1]

- (b) Photochromic glass used for sunglasses contains AgCl crystals which are added while the glass is in molten state. The glass darkens when exposed to bright light and the following reaction is involved.




- (i) Suggest why the glass darkens when exposed to bright light. [1]

The glass darkens significantly within about a minute of exposure to bright light but takes a longer time to clear when there is less light. Small amount of CuCl crystals are often added to the glass to speed up the reverse process. No elemental copper is involved in the process.



- (ii) Explain why the copper(I) ions can be described as a *homogeneous catalyst*. [2]
- (iii) State the property, typical of transition metals, which allows copper(I) ions to behave as a homogenous catalyst in the reverse reaction. Include relevant chemical equations to support your answer. [3]
- (c) Microorganisms synthesise and secrete organic molecules called siderophores to increase the total concentration of available iron in the surrounding medium. Ferrichrome is a siderophore produced by fungi.




Ferrichrome

- (i) Ferrichrome binds to iron(III) ions via its oxygen atoms. This process facilitates the transportation of iron(III) ions into the interior of a cell.
Suggest what bonds are formed during this process and why iron(III) does not bind to nitrogen atoms in ferrichrome. [2]
- (ii) Draw the structures of the products formed when ferrichrome is heated with aqueous sodium hydroxide. [3]


[Total: 20]

- 2 Ibuprofen is a nonsteroidal anti-inflammatory drug. It works by reducing hormones that cause inflammation and pain in the body. Ibuprofen is used to reduce fever and treat pain or inflammation.


The conventional synthesis of ibuprofen from isobutylbenzene is shown below. Step 4 of the synthesis resembles the reaction between a carbonyl compound and Brady's reagent.



- (a) Name the types of reaction that are occurring during steps 4 and 5. [2]
- (b) Suggest the reagents and conditions for steps 1, 4 and 6. [3]
- (c) Draw the structures of the stereoisomers of the product formed from step 4 and state the type of stereoisomerism exhibited. [2]
- (d) Describe a simple chemical test to distinguish between the two compounds, **X** and **Y** obtained from synthesis above. [2]



Compound X



Compound Y

- (e) Potassium chloride and potassium iodide can be distinguished by treating the compounds separately with concentrated sulfuric acid.

- (i) Potassium chloride and concentrated sulfuric acid reacts in an equimolar ratio to produce white fumes of hydrogen chloride.


Write an equation for the reaction of potassium chloride with concentrated sulfuric acid. [1]

- (ii) Potassium iodide reacts with concentrated sulfuric acid in a similar manner. However, the white fumes of hydrogen iodide would further react with concentrated sulfuric acid to produce violet fumes and hydrogen sulfide gas.

Suggest an explanation for the difference in reactions, and write equations for the above observations. [2]

- (f) When aqueous potassium salts of a dicarboxylic acid are electrolysed using inert platinum electrodes, alkenes are formed at the anode.

- (i) The electrolysis of aqueous potassium succinate gives ethene at the anode.




Potassium Succinate

Write an ion-electron half equation for the oxidation of succinate at the anode. [1]

- (ii) Write an equation for the reaction at the cathode and for the overall reaction. [2]

- (iii) The mechanism of the reaction at the anode involves three steps:


Step 1: There is an initial loss of 2 electrons on succinate ion to form



Step 2: This is followed by decarboxylation which involves the homolytic breaking of two C–C bonds, giving a radical intermediate $\cdot\text{CH}_2\text{CH}_2\cdot$

Step 3: The third step involves forming a covalent bond, producing ethene as the product.



Use the information given above to draw the mechanism for Steps 2 and 3.



You are advised to use structural formulae for all species, such as $\cdot \text{O} - \overset{\text{O}}{\parallel} \text{C} - \underset{\text{H}}{\underset{|}{\text{C}}} - \underset{\text{H}}{\underset{|}{\text{C}}} - \overset{\text{O}}{\parallel} \text{C} - \text{O} \cdot$, so that it is clear which bonds are broken and which are formed.

Represent with a half arrow ($\curvearrowright$) for movement of a single unpaired electron and indicate any unpaired electrons by a dot ($\cdot$). [2]

- (g) The strength of a carboxylic acid depends on its structure. An example of this is the comparison of succinic acid and its derivatives.

Acid	pK_{a1}
	4.21
	-


Explain if pK_a of chlorosuccinic acid is lower or higher than the pK_a of succinic acid. [1]

[Total: 18]

BLANK PAGE


- 3 On 28 November 2018, an explosion erupted near the Hebei Shenghua Chemical Industrial plant, killing at least 23 people and injuring many more. It was reported that the explosion originated from a truck transporting ethyne gas, C_2H_2 , which then set off a chain reaction that engulfed at least 50 other vehicles.

When mixed appropriately with air, C_2H_2 explodes upon ignition as follows.



This explosion may have generated an estimated 20.8 million kilojoules of energy.

- (a) Describe the hybridisation of the orbitals in, and the bonds between, the carbon atoms within an ethyne molecule. [3]
- (b) (i) Using the bond energies in the *Data Booklet*, calculate the enthalpy change of combustion of ethyne. [2]
- (ii) Using your answer in (b)(i), calculate the volume of ethyne gas transported by the truck at 28°C and 1 bar. [2]
- (c) One of the ways that ethyne gas can be produced is via the reaction of calcium carbide, CaC_2 , with water, producing solid calcium hydroxide as a by-product.
- (i) Using the following information and relevant data from the *Data Booklet*, construct an energy cycle to calculate the lattice energy of solid calcium hydroxide. [3]



standard enthalpy change of atomisation of Ca (s) = + 178 kJ mol^{-1}


standard enthalpy change of formation of $\text{OH}^{-}(\text{g})$ = – 230 kJ mol^{-1}

standard enthalpy change of formation of $\text{Ca}(\text{OH})_2(\text{s})$ = – 985 kJ mol^{-1}

- (ii) Suggest, with reasoning, how the magnitude of the lattice energy of $\text{Ca}(\text{OH})_2$ compares with that of calcium carbide, CaC_2 , given the following data. [2]

Ion	Radius (pm)
OH^{-}	133
C_2^{2-}	118


- (iii) The value of $\Delta G^\ominus$ for the reaction of calcium carbide with water is -148 kJ mol^{-1} .



Calculate $\Delta S^\ominus$ in $\text{J mol}^{-1} \text{ K}^{-1}$ to one decimal place for the reaction. Explain the significance of its sign with reference to the equation. [2]

- (d) Similar to calcium carbide, sodium borohydride, NaBH_4 , can be used in fuel cells.

Direct borohydride fuel cells (DBFCs), are fuel cells which are directly fed by NaBH_4 as the fuel and oxygen as the oxidant. The electrolyte used is potassium hydroxide. Sodium metaborate, NaBO_2 , and water are formed as the only products during the discharging process.



- (i) Deduce the ion-electron half-equations for the anode and cathode and hence write the overall cell reaction during discharging. [2]
- (ii) A typical DBFC fuel cell generates about 1.64 V. Using the *Data Booklet*, calculate the $E^\ominus$ involving the $\text{NaBO}_2/\text{NaBH}_4$ half-cell. [1]
- (iii) Use the half-equations you have written in (d)(i) to calculate the value of $\Delta G^\ominus$ and comment on the significance of $\Delta G^\ominus$. [1]
- (iv) Explain qualitatively what happens to the cell potential, E_{cell} when a small amount of propanal contaminant is accidentally added to the anode half-cell.
The contaminant does not take part in the cell reaction. [2]
- (v) In a particular DBFC, a current of $1.35 \times 10^{-2} \text{ A cm}^{-3}$ was passed through the circuit for 95 minutes.
Calculate the mass of NaBO_2 that was produced per 25 cm^3 of solution. [2]

[Total: 22]

Section B

Answer **one** question from this section.

- 4 Chlorine-containing compounds have many useful applications. One such example is the use of chloric(I) acid, HClO , to disinfect water in swimming pools.

HClO dissociates in water as shown below.



- (a) (i) Write an expression for K_a of HClO . [1]

- (ii) A disinfectant solution was prepared by dissolving 10 g of HClO and x g of NaClO in 100 cm^3 of water.

For effective disinfection, a pH of 7.35 was needed to reach the optimal $[\text{ClO}^-]:[\text{HClO}]$ ratio.


Given that the $\text{p}K_a$ of HClO is 7.55, determine the optimal $[\text{ClO}^-]:[\text{HClO}]$ ratio and hence the mass of NaClO , x . [2]

- (iii) A worker accidentally poured some alkaline solution into the disinfectant solution formed in (a)(ii) but found that the pH was still about 7.35.

Write an equation to explain this. [1]


- (b) Hydrogen chloride is another chlorine-containing compound.

Alkyl chlorides can be synthesised from alcohols using hydrogen chloride. However, acyl chlorides **cannot** be prepared from carboxylic acids in the same way. One key factor is due to the stronger C-O bond present in carboxylic acids.




Suggest an explanation for the stronger C-O bond in carboxylic acids. [1]

- (c) Acetanilide is an analgesic and can be formed from ethanoyl chloride and phenylamine.



Instead of ethanoyl chloride, ethanoic anhydride is more commonly used in the laboratory to form acetanilide in a similar manner.



- (i) Suggest the other organic product that is produced together with acetanilide. [1]

The experimental procedure for the laboratory synthesis of acetanilide using ethanoic anhydride is given below.

1. Mix phenylamine and hydrochloric acid in a beaker.
2. Stir the mixture until a clear solution is obtained.
3. To the clear solution, add ethanoic anhydride and immediately add a solution of sodium ethanoate.
4. Stir the mixture vigorously.
5. Cool the solution in an ice water bath.
6. Filter the acetanilide formed. Wash with cold water and crystallise using a mixture of water and methanol.



- (ii) In step 1, HCl was added to increase the solubility of phenylamine in water. Explain. [1]

- (iii) Sodium ethanoate is essential to ensure phenylamine is present in step 3 to react with ethanoic anhydride.

Suggest the role of sodium ethanoate. [1]

- (iv) Suggest how you could check that the acetanilide obtained in step 6 is pure. [1]

- (d) Aspirin is another analgesic and it can be synthesised from salicylic acid. The pK_a values of salicylic acid and its isomer are given below:

Acid	pK_{a1}	pK_{a2}
 Salicylic acid	3.0	13.4
 4-hydroxybenzoic acid	4.1	9.7

Explain the following:

- (i) pK_{a1} of salicylic acid is smaller than pK_{a1} of 4-hydroxybenzoic acid. [1]
- (ii) pK_{a2} is larger than pK_{a1} of 4-hydroxybenzoic acid. [1]
- (e) When compound **A**, $C_{10}H_{11}NO$, is oxidised with acidified manganate(VII) ions, $[C_7H_8NO_2]^+$ and compound **B**, $C_3H_4O_3$, are formed. Compound **B** produces effervescence when Na_2CO_3 is added to it and gives an orange precipitate with 2,4-dinitrophenylhydrazine reagent.

Compound **A** reacts readily with 4 moles of $Br_2(aq)$ to give compound **C**, $C_{10}H_9NO_2Br_4$.

Compound **A** also gives an orange precipitate with 2,4-dinitrophenylhydrazine reagent and reacts with hot alkaline aqueous iodine to give compound **D**, $C_9H_8NO_2Na$.


Suggest the structures of compounds **A**, **B**, **C** and **D**. Explain the reactions that occur. [9]

[Total: 20]


BLANK PAGE

- 5 (a) Hydrogen sulfide gas, H_2S , is a reagent used in qualitative analysis. When H_2S is passed into a solution containing metal cations, metal sulfides are precipitated.

In aqueous medium, H_2S dissociates partially in a step-wise manner to give two equilibrium reactions:



The relationship between the H^+ ion concentration and S^{2-} ion concentration, as a result of H_2S dissociation, is shown in the graph below, where $\text{pS} = -\log_{10}[\text{S}^{2-}]$.



Solution **X** contains $0.1 \text{ mol dm}^{-3} \text{ CuSO}_4$ and $0.05 \text{ mol dm}^{-3} \text{ MnSO}_4$. Hydrogen sulfide gas is bubbled into **X** until saturation and the pH is maintained at 2.0.

The value of the solubility product of CuS and MnS are 1.0×10^{-44} and 1.4×10^{-15} , respectively.

- (i) With reference to CuS , define the term *solubility product*. [1]
- (ii) Calculate the minimum concentration of sulfide ions needed to precipitate CuS . [1]
- (iii) The minimum concentration of sulfide ions needed to precipitate MnS is $2.80 \times 10^{-14} \text{ mol dm}^{-3}$.

Using the graph above and your answer in (a)(ii), explain whether it is possible to separate copper and manganese ions in **X** at $\text{pH} = 2$. [2]

- (iv) Determine the maximum pH for solution **X** before separation of copper and manganese ions is no longer possible. [1]

- (b) When water ligands in a hydrated metal ion are substituted by other ligands, the equilibrium constant for the reaction is referred to as the stability constant, K_{stab} , of the new complex. The higher the value of K_{stab} , the more likely the complex will be formed.


The stability constants for 3 reactions are given below.

Reaction	Colour of complex formed	Value of $\log_{10}(K_{\text{stab}})$
$\text{Fe}^{3+}(\text{aq}) + 6\text{CN}^{-}(\text{aq}) \rightleftharpoons [\text{Fe}(\text{CN})_6]^{3-}(\text{aq})$	Yellow	31.0
$\text{Fe}^{3+}(\text{aq}) + \text{EDTA}^{4-}(\text{aq}) \rightleftharpoons [\text{Fe}(\text{EDTA})]^{-}(\text{aq})$	Brown	25.1
$\text{Fe}^{3+}(\text{aq}) + \text{SCN}^{-}(\text{aq}) \rightleftharpoons [\text{Fe}(\text{SCN})]^{2+}(\text{aq})$	Red	2.1

Describe the colour change(s) of the solution when excess NaSCN is added to $[\text{Fe}(\text{EDTA})]^{-}$, followed by an excess of KCN(aq).


Explain the observation, with reference to the table above. [2]

- (c) A solution of $\text{Pb}(\text{NO}_3)_2$ is subjected to the following reactions.



- (i) Suggest the identity of white solid **A** and complex anion present in colourless solution **B**. [2]
- (ii) Colourless gas **C** is obtained from the reaction of solid PbS and concentrated HNO_3 , which is readily oxidised in air to form a brown acidic gas. Suggest the identity of **C**. [1]
- (d) Thiols are sulfur-containing compounds that can be prepared from alkyl halides and the hydrosulfide nucleophile, SH^{-} .

Ethanethiol is prepared from chloroethane using NaSH as shown below.




- (i) Suggest a synthesis of propanethiol starting from propylethanoate. You should state the reagents and conditions for each step, and show clearly the structure of any intermediate compounds. [3]

$\text{CH}_3\text{CH}_2\text{SH}$ can react with a base, NaH , to give $\text{CH}_3\text{CH}_2\text{S}^-$.



$\text{CH}_3\text{CH}_2\text{S}^-$ further reacts with chloroethane, to yield diethyl sulfide



Diethyl sulfide


(ii) Suggest why $\text{CH}_3\text{CH}_2\text{S}^-$, is a better nucleophile than $\text{CH}_3\text{CH}_2\text{O}^-$. [1]

(iii) Diethyl sulfide is a good nucleophile.

In Reaction 2, a side product, $\text{C}_6\text{H}_{15}\text{SCl}$, is formed when chloroethane is used in excess. $\text{C}_6\text{H}_{15}\text{SCl}$ gives a white precipitate with aqueous silver nitrate.

Suggest the structural formula of the side product $\text{C}_6\text{H}_{15}\text{SCl}$. [1]

(e) The millipede *Apheeloria corrugate* protects itself by secreting the cyanohydrin mandelonitrile and an enzyme that decompose mandelonitrile into benzaldehyde and HCN .



(i) State the type of reaction for the formation of Mandelonitrile from benzaldehyde. [1]

(ii) Suggest a chemical test that will give a positive test for Mandelonitrile but not for benzaldehyde. [2]

(iii) A student made the following statement, "benzaldehyde can be directly synthesized from phenylmethanol, $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$, but not from benzoic acid."

Based on your knowledge of the reagents and conditions for each reaction, deduce if the statement is true. [2]

[Total: 20]