


2022 Prelim A Maths Paper 1 Marking Scheme

Qn	Solution	Marks	Remarks
1	$y = x^2 + 3x - 8 \text{ ----- (1)}$ $y = \frac{3}{2}x - \frac{7}{2} \text{ ----- (2)}$ $(1) = (2)$ $x^2 + 3x - 8 = \frac{3}{2}x - \frac{7}{2}$ $2x^2 + 3x - 9 = 0$ $(2x - 3)(x + 3) = 0$ $x = \frac{3}{2} \text{ or } x = -3$ $y = -\frac{5}{4} \text{ or } y = -8$ Coord of A and B are $\left(\frac{3}{2}, -\frac{5}{4}\right)$ & $(-3, -8)$. Distance btw A and B $= \sqrt{(1.5 + 3)^2 + (-1.25 + 8)^2}$ $= 8.11 \text{ units}$	[M1] [M1] [M1] [A1]	Accept exact surd $\frac{\sqrt{1053}}{4}$
2	$\frac{7x+8}{(2x+1)(x-1)^2} = \frac{A}{2x+1} + \frac{B}{x-1} + \frac{C}{(x-1)^2}$ $7x - 8 = A(x-1)^2 + B(2x+1)(x-1) + C(2x+1)$ Let $x = 1$, $15 = 3C$, $C = 5$ Let $x = -\frac{1}{2}$, $\frac{9}{2} = A\left(-\frac{3}{2}\right)^2$, $A = 2$ Let $x = 0$, $8 = 2(-1)^2 + B(1)(-1) + 5(1)$, $B = -1$ $\frac{7x+8}{(2x+1)(x-1)^2} = \frac{2}{2x+1} - \frac{1}{x-1} + \frac{5}{(x-1)^2}$	[M1] [M1] [A3]	

(b)	$f(x) = \int (3x^2 + 2x + 10) dx$ $= x^3 + x^2 + 10x + d$ At $(-1, 10)$, $10 = (-1)^3 + (-1)^2 + 10(-1) + d$ $d = 20$ $f(x) = x^3 + x^2 + 10x + 20$	[M1] [A1]	
(c)	For $y = f(x)$ to have stationary points, set $f'(x) = 0$. $3x^2 + 2x + 10 = 0$ $b^2 - 4ac = 2^2 - 4(3)(10) = -116 < 0$ $f'(x) = 0$ has no real solution, ie no stationary points.	[M1] [M1] [A1]	Solve equation to show no real roots accepted No marks if no real roots is not mentioned
7(a)	$x = 4.5, \quad V = \frac{1}{3} \pi (4.5)^2 (18 - 4.5)$ $= \frac{729}{8} \pi$ $\frac{dV}{dt} = \frac{\frac{729}{8} \pi}{9} = \frac{81}{8} \pi \text{ cm/s (or } 10\frac{1}{8} \pi \text{ cm/s)}$	[M1] [M1.A1]	Accept 10.125π or 31.8 cm/s
(b)	$V = \frac{\pi}{3} (18x^2 - x^3)$ $\frac{dV}{dx} = \frac{\pi}{3} (36x - 3x^2)$ At $x = 4.5$, $\frac{dV}{dx} = \frac{\pi}{3} (36(4.5) - 3(4.5)^2)$ $= \frac{135}{4} \pi$ Using $\frac{dV}{dt} = \frac{dV}{dx} \cdot \frac{dx}{dt}$ $\frac{dx}{dt} = \frac{\frac{81}{8} \pi}{\frac{135}{4} \pi} = 0.3$ The water level is rising at a rate of 0.3 cm/s	[M1] [M1] [M1.A1]	

8(a)	$\sin^3 x + \cos^3 x$ $= (\sin x + \cos x)(\sin^2 x - \sin x \cos x + \cos^2 x)$	[B1]	
(b)	$LHS = \frac{\sin^3 x + \cos^3 x}{\sin x + \cos x}$ $= \frac{(\sin x + \cos x)(\sin^2 x - \sin x \cos x + \cos^2 x)}{\sin x + \cos x}$ $= \sin^2 x - \frac{1}{2}(2 \sin x \cos x) + \cos^2 x$ $= 1 - \frac{1}{2} \sin 2x = RHS$	[M1] [A1, A1]	No mark awarded if student do not show $\sin x \cos x = \frac{1}{2}(2 \sin x \cos x)$
(c)	$1 - \frac{1}{2} \sin 2x = \frac{5}{4}$ $\frac{1}{2} \sin 2x = -\frac{1}{4}$ $\sin 2x = -\frac{1}{2}$ Basic angle, $\alpha = 30^\circ$ $2x = 210^\circ, 330^\circ, 570^\circ, 690^\circ$ $x = 105^\circ, 165^\circ, 285^\circ, 345^\circ$	[M1] [M1] [M1] [A1]	
9(a)	$\angle ACB = \angle ATC$ (given) $\angle BAC = \angle CAT$ (common) $\therefore \triangle ABC \text{ \& } \triangle ACT$ are similar (AA Test)	[M1] [A1]	Either statement
(b)	Since $\triangle ABC \text{ \& } \triangle ACT$ are similar , $\frac{AB}{AC} = \frac{AC}{AT}$ $AC^2 = (AB)(AT)$ $= (AT + TB)(AT)$ $= AT^2 + AT \times TB$ $AC^2 - AT^2 = AT \times TB \text{ (shown)}$	[M1] [M1] [A1]	
(c)	$\angle BAX = \angle ACB$ (alt. segment thm) $\& \angle ACB = \angle ATC$ (given) $\therefore \angle BAX = \angle ATC$ By the alternate angle property, SC and XY are parallel.	[M1] [M1] [A1]	

10(a)	$m_{AB} = \frac{p-10}{3}$ $m_{CB} = \frac{6-p}{1}$ Since $\angle ABO = \angle CBO, m_{AB} = -m_{CB}$ $\frac{p-10}{3} = -(6-p)$ $p-10 = -18+3p$ $-2p = -8$ $p = 4 \quad (\text{shown})$	[M1] [M1] [A1]	Accept method involving $\tan \theta$
(b)	$m_{AB} = -2$ $m_{AD} = \frac{1}{2}$ Equation of line AD is $y-10 = \frac{1}{2}(x+3)$ $y = \frac{1}{2}x + \frac{23}{2} \quad (\text{shown})$ $CD \parallel AB, m_{CD} = -2$ Equation of line CD is $y-6 = -2(x-1)$ $y = -2x+8$ At D, $\frac{1}{2}x + \frac{23}{2} = -2x+8$ $\frac{5}{2}x = -\frac{7}{2}$ $x = -\frac{7}{5}, y = -2\left(-\frac{7}{5}\right) + 8 = \frac{54}{5}$ Coord of D is $\left(-\frac{7}{5}, \frac{54}{5}\right)$	[M1] [A1] [M1] [M1]	Accept (-1.4, 10.8)

(c)	Area of $ABCD$ $= \frac{1}{2} \begin{vmatrix} 0 & 1 & -1.4 & -3 & 0 \\ 4 & 6 & 10.8 & 10 & 4 \end{vmatrix}$ $= \frac{1}{2} [(0+10.8-14-12)-(4-8.4-32.4)]$ $= 10.8 \text{ units}^2$	[M1] [A1]	
11(a)	$x^2 + 4x + 3 = (x+3)(x+1)$ By Factor Theorem, $P(-3) = 0$ & $P(-1) = 0$ $5(-3)^3 + a(-3)^2 + 3 + b = 0$ $9a + b = 132 \quad \text{-----(1)}$ $5(-1)^3 + a(-1)^2 + 1 + b = 0$ $a + b = 4 \quad \text{-----(2)}$ (1) – (2): $8a = 128$ $a = 16, \quad b = -12$	[M1] [M1] [M1] [A1]	*overall minus 1 mark if any expression is not set to 0
(b)	$5x^3 + 16x^2 - x - 12 = (x+3)(x+1)(px+q)$ By comparison, $p = 5$ & $q = -4$ $P(x) = 0$ $(x+3)(x+1)(5x-4) = 0$ $x = -3$ or -1 or $\frac{4}{5}$	[M1] [M1] [A1]	
(c)	$5(x^2)^3 + a(x^2)^2 - x^2 + b = 0 \quad \text{---- (1)}$ Let $u = x^2$, (1) becomes $5u^3 + au^2 - u + b = 0$ From (b), $u = -3(\text{rej})$ or $-1(\text{rej})$ or $\frac{4}{5}$ $\therefore x^2 = \frac{4}{5}$ $x = \frac{2}{\sqrt{5}}$ or $-\frac{2}{\sqrt{5}}$	[M1] [A1]	Accept $\pm \frac{2\sqrt{5}}{5}$

12(a)	$y = \frac{7}{6x+1} = 7(6x+1)^{-1}$ $\frac{dy}{dx} = -7(6x+1)^{-2} (6)$ $= -\frac{42}{(6x+1)^2}$ At K, $x = 1$, $\frac{dy}{dx} = -\frac{42}{7^2} = -\frac{6}{7}$ Equation of tangent at K is $y - 1 = -\frac{6}{7}(x - 1)$ $y = -\frac{6}{7}x + \frac{13}{7}$ Coord of B is $\left(0, \frac{13}{7}\right)$	[M1] [M1] [M1] [M1] [A1]	
(b)	Area of the shaded region = Area under curve – Area of trapezium $= \int_0^1 \frac{7}{6x+1} dx - \frac{1}{2}(1)\left(1 + \frac{13}{7}\right)$ $= 7 \left[\frac{1}{6} \ln(6x+1) \right]_0^1 - \frac{10}{7}$ $= \frac{7}{6}(\ln 7 - \ln 1) - \frac{10}{7}$ $= 0.842 \text{ units}^2$	[M1,,M1] [M1] [M1] [A1]	
13(a)	$(\ln x)^2 + \frac{2}{\frac{\ln e}{\ln x}} = 3$ $(\ln x)^2 + 2 \ln x - 3 = 0$ Let $u = \ln x$ $u^2 + 2u - 3 = 0$ $(u+3)(u-1) = 0$ $u = -3 \text{ or } u = 1$ $\ln x = -3 \text{ or } \ln x = 1$ $x = e^{-3} = \frac{1}{e^3} \text{ or } x = e$	[M1] [M1] [M1] [A1]	Accept $x = 0.0498$ or $x = 2.72$

(b)	$\lg\left(\frac{p}{2q}\right) = \lg(p + 2q)$ $\frac{p}{2q} = p + 2q$ (i) $p = 2pq + 4q^2$ $p(1 - 2q) = 4q^2$ $p = \frac{4q^2}{1 - 2q}$	[M1]	
	(ii) Range of p is $p > 0$ $\therefore \frac{4q^2}{1 - 2q} > 0$ Since $q > 0$ for $\lg 2q$ to be defined, $4q^2 > 0 \text{ \& } 1 - 2q > 0$ $q < \frac{1}{2}$ $\therefore 0 < q < \frac{1}{2} \text{ (shown)}$	[B1]	
(c)	By observing the shape of the curve, a logarithmic function, ie equation (B) $y = a \ln x + b$ is a suitable model since the rate of growth of the head circumference gets much slower when the baby gets older over the months.	[B1] [B1]	-for correct model -for correct reasoning