



TAMPINES MERIDIAN JUNIOR COLLEGE

JC2 PRELIMINARY EXAMINATION

H1 CHEMISTRY

8873/01

Paper 1 Multiple Choice

19 September 2024

1 hour

Additional materials: Multiple Choice Answer Sheet
Data Booklet

READ THESE INSTRUCTIONS FIRST

Write in soft pencil.

Write your name, class and register number on the Answer Sheet in the spaces provided.

There are **thirty** questions in this section. Answer **all** questions. For each question, there are four possible answers labelled **A, B, C** and **D**.

Choose the **one** you consider correct and record your choice in soft pencil on the separate Answer Sheet.

Read very carefully the instructions on the use of Answer Sheet.

You are advised to fill in the Answer Sheet as you go along; no additional time will be given for the transfer of answers once the examination has ended.

Use of Answer Sheet

Ensure you have written your name, class register number and class on the Answer Sheet.

Use a **2B** pencil to shade your answers on the Answer Sheet; erase any mistakes cleanly. Multiple shaded answers to a question will not be accepted.

For shading of class register number on the **Answer Sheet**, please follow the given examples:

If your register number is **1**, then shade **01** in the index number column.

If your register number is **21**, then shade **21** in the index number column.

- 1 *Use of the Data Booklet is relevant to this question.*

A stream of gaseous $^{40}_{20}\text{Ca}^{2+}$ charged ions was passed through an electric field between two oppositely charged plates and the Ca^{2+} ions were deflected at an angle of $+15.0^\circ$.

Under the same electric field, a stream of unknown gaseous X^{3-} ion was deflected at an angle of -29.0° . What is a possible atomic number of **X**?

- A** 15 **B** 24 **C** 31 **D** 34

- 2 Which of these statements correctly describes an electron shell with the principal quantum shell number $n = 2$?

- 1 A total of 10 electrons can be accommodated in this shell.
- 2 An orbital from this shell must be dumb-bell in shape.
- 3 The energy level of the orbitals in this shell is higher than that in $n = 1$.

- A** 1, 2 and 3 **B** 1 and 2 only **C** 2 and 3 only **D** 3 only

- 3 *Use of the Data Booklet is relevant to this question.*

The first ionisation energy of beryllium is greater than the first ionisation energy of boron.

Which factor explains this?

- A** charge on the nucleus
B distance from the nucleus to the outer electrons
C removal of electron is from a lower energy orbital
D repulsion between electrons in an orbital

- 4 Which of the following does **not** contain hydrogen bonding?

- A** $\text{NH}_3(\text{l})$ **B** $\text{NH}_4\text{Cl}(\text{l})$ **C** $(\text{CH}_3)_2\text{NH}(\text{l})$ **D** $\text{CH}_3\text{CONH}_2(\text{l})$

- 5 The CO_2 molecule is linear.

What is the number of σ and π bonds present in the molecule?

- | | σ | π |
|----------|----------|-------|
| A | 1 | 3 |
| B | 2 | 2 |
| C | 3 | 1 |
| D | 4 | 0 |



- 6 The table identifies the shape and polarity of four molecules. Which row is correct?

	molecule	molecular shape	polarity
A	bromine trifluoride	bent	polar
B	phosphorus trichloride	trigonal pyramidal	polar
C	sulfur dichloride	linear	non-polar
D	trifluoromethane	tetrahedral	non-polar

- 7 The table shows four species with their corresponding physical properties. Which species does **not** correspond to its description of physical properties?

	species	physical properties
A	copper	high melting point, conducts electricity when solid and when molten
B	silicon dioxide	high melting point, does not conduct electricity in any state
C	aluminium bromide	high melting point, conducts electricity when molten but not when solid
D	phosphorus pentachloride	low melting point, does not conduct electricity in any state


- 8 Which of the following is **not** an acid-conjugate base pair?

- A** HCl and NaCl
- B** Na_2HPO_4 and H_3PO_4
- C** H_2O_2 and HO_2^-
- D** $\text{CH}_3\text{CH}_2\text{NH}_2$ and $\text{CH}_3\text{CH}_2\text{NH}_3\text{Cl}$

- 9 Which one of the following acid solutions can be used to give an effective buffer solution by partial neutralisation with aqueous NaOH?

A HCOOH
 B HCl
 C HI
 D HNO_3

- 10 The graph shown can be plotted for the titration of aqueous ethanoic acid with potassium hydroxide.



Which species are present in the solution at point X?

- 1 ethanoic acid
 2 potassium hydroxide
 3 potassium ethanoate


A 1 and 2 only B 1 and 3 only C 2 and 3 only D 3 only

- 11 Which pair of statements is correct for Group 1 elements and their reducing strength?

	reducing strength	reason
A	decreases down the Group	valence electron more easily lost
B	decreases down the Group	valence electron more easily gained
C	increases down the Group	valence electron more easily lost
D	increases down the Group	valence electron more easily gained

12 Use of the Data Booklet is relevant to this question.

Which of the following shows the correct trends when the melting points of the elements, Mg, Al, Si and P are plotted against their atomic radii?



13 Which statement is correct about the halogens (X_2) chlorine, bromine and iodine as well as their corresponding hydrogen halides (HX)?

- A The colour intensity of the halogens decreases from Cl_2 to I_2 .
- B The oxidising power of the halogens decreases from Cl_2 to I_2 .
- C The thermal stability of HX increases from HCl to HI.
- D The volatility of HX increases from HCl to HI.

14 1 dm^3 of a gaseous hydrocarbon was mixed with excess oxygen. When the mixture undergoes complete combustion, the volume contracted by 2 dm^3 . When the resultant mixture was passed through concentrated sodium hydroxide, the gas volume further decreased by 2 dm^3 . All volumes were measured at room temperature and pressure.

What is the empirical formula of the hydrocarbon?

- A CH
- B CH_2
- C C_2H_3
- D C_3H_4




- 15 In this question, only V and Zn undergo changes in oxidation numbers.

Vanadium oxide, V_2O_5 , reacts with zinc in the presence of acidic medium. Zn is oxidised to Zn^{2+} and 3 moles of Zn react with 1 mole of V_2O_5 .

What is the final oxidation number of V in the vanadium containing product of this reaction?

- A** 0 **B** +1 **C** +2 **D** +3

- 16 Chromates such as sodium chromate, Na_2CrO_4 , are used in chrome plating to protect metals and can be manufactured by heating iron(II) chromite, $FeCr_2O_4$, with sodium carbonate in air.



Which species are oxidised and reduced in this process?

	species oxidised	species reduced
A	chromium and iron	carbonate ion and oxygen
B	chromium and iron	oxygen only
C	iron only	carbonate ion only
D	chromium only	carbonate ion and oxygen

- 17 Substances **X** and **Y** react to form product **Z**.



The reaction was studied and the following data were obtained.


experiment	[X] / mol dm ⁻³	[Y] / mol dm ⁻³	initial rate / mol dm ⁻³ min ⁻¹
1	0.10	0.10	0.020
2	0.20	0.10	0.080
3	0.15	0.23	0.045

What is the order of reaction with respect to **X** and **Y**?

- A** First order with respect to **X** and zero order with respect to **Y**.
B Second order with respect to **X** and first order with respect to **Y**.
C First order with respect to **X** and first order with respect to **Y**.
D Second order with respect to **X** and zero order with respect to **Y**.



- 18 The graph shows the results of decomposition of compound A.



Which conclusions can be drawn from the graph?

- 1 The half-life of the decomposition of A is constant.
- 2 The rate of the reaction is independent of the concentration of A.
- 3 Units of rate constant, k , is $\text{mol dm}^{-3} \text{min}^{-1}$.

A 1, 2 and 3 **B** 1 and 2 only **C** 2 and 3 only **D** 1 only

- 19 A radioactive isotope follows first order kinetics in its decay.

If the rate of decay of a radioactive isotope decreases from 400 counts per minute to 25 counts per minute after 24 hours, what is its half-life?

- A** 3 hours
B 4 hours
C 6 hours
D 12 hours

- 20 Which equation best represents the enthalpy change of formation of iron(III) oxide, Fe_2O_3 ?

- A** $2\text{Fe(l)} + 3\text{O(g)} \rightarrow \text{Fe}_2\text{O}_3\text{(s)}$
B $4\text{Fe(s)} + 3\text{O}_2\text{(g)} \rightarrow 2\text{Fe}_2\text{O}_3\text{(s)}$
C $2\text{Fe(s)} + 3\text{O(g)} \rightarrow \text{Fe}_2\text{O}_3\text{(s)}$
D $2\text{Fe(s)} + \frac{3}{2}\text{O}_2\text{(g)} \rightarrow \text{Fe}_2\text{O}_3\text{(s)}$


- 21** The enthalpy change of solution of a solid ionic compound is the energy change when one mole of the substance is completely dissolved in a solvent to form aqueous ions.

The enthalpy change of solution is dependent on:

- lattice energy of the ionic compound,
- hydration energy of the gaseous ions to aqueous ions.

	/ kJ mol^{-1}
Lattice energy of AlF_3	-6083
Enthalpy change of hydration of Al^{3+}	-4750
Enthalpy change of hydration of F^-	-473

Using the energy level diagram and the data provided, determine the enthalpy change of solution of aluminium fluoride, $\text{AlF}_3(\text{s})$.




- A** -86 kJ mol^{-1}
B -860 kJ mol^{-1}
C $-11300 \text{ kJ mol}^{-1}$
D $-12250 \text{ kJ mol}^{-1}$

- 22 The following enthalpy changes of the following substances are provided.

	/ kJ mol ⁻¹
Standard enthalpy change of combustion of graphite	-394
Standard enthalpy change of formation of water	-286
Standard enthalpy change of formation of liquid ethanol	-277



Calculate the value of standard enthalpy change of combustion of liquid ethanol.

- A -403 kJ mol⁻¹
 B -957 kJ mol⁻¹
 C -1369 kJ mol⁻¹
 D -1865 kJ mol⁻¹
- 23 Consider the following equilibrium reaction:



Which of the following changes will decrease the equilibrium yield of HCl?


- A adding more H₂(g)
 B adding a suitable catalyst
 C decreasing the volume of the system
 D increasing the temperature
- 24 The graph below shows how the percentage of SO₃(g) in the equilibrium mixture of Contact Process varies with temperature at different pressures, p₁ and p₂.



Which of the following is correct?

- | | ΔH_r | pressure |
|---|--------------|-------------|
| A | exothermic | $p_1 < p_2$ |
| B | exothermic | $p_1 > p_2$ |
| C | endothermic | $p_1 < p_2$ |
| D | endothermic | $p_1 > p_2$ |

- 25** A mixture of hydrogen and iodine will form an equilibrium mixture as shown by the equation.



The table gives the concentration of each substance in a mixture at equilibrium under certain conditions.

substance	concentration / mol dm ⁻³
H ₂	0.6
I ₂	0.2
HI	0.4


What is the numerical value of K_c under these conditions?

- A** 0.30
B 0.75
C 1.33
D 3.33
- 26** Properties of graphene make it ideal for making the next-generation optoelectronics and optical communications systems, particularly in making touchscreens, liquid crystal displays (LCD) and organic light emitting diodes (OLEDs).

Which property of graphene is the least essential for its use in touchscreen tablets?


- A** good electrical conductivity
B high optical transparency
C high tensile strength
D light weight

- 27 When exposed to chlorine gas under UV light, 2-methylbutane formed different mono-chlorinated products.



What is the ratio of the mono-chlorinated products formed?

- A 7 : 5
 B 6 : 5 : 1
 C 9 : 2 : 1
 D 6 : 3 : 2 : 1
- 28 What is the total number of constitutional isomers, including ring structures, that has the molecular formula C_4H_8 ?
- A 3 B 4 C 5 D 6
- 29 Polybutylene succinate (PBS) is a biodegradable polymer which found itself useful in many applications like disposable food packaging and medicine capsules.




PBS repeating unit

The PBS molecule is made up of two different monomer units. Which of these can be the monomers?

- A
- B
- C
- D

- 30 The flexibility and strength of 4 polymers Q to T are measured.

They are first made into sheets. These sheets are supported at opposite ends and masses are added to the centre of the sheets. As the masses are added, the amount the polymer bends, in degrees, is measured until the polymer breaks.



Which conclusions are true?

- 1 Polymer Q is most suitable for making chairs.
- 2 Polymer R is likely a thermoplastic.
- 3 Polymer S is most suitable for making machinery parts.
- 4 Polymer T is likely a thermosetting plastic.

A 1, 2, 3 and 4 **B** 2, 3 and 4 only **C** 1 and 3 only **D** 2 and 4 only