



# TAMPINES MERIDIAN JUNIOR COLLEGE

## JC2 PRELIMINARY EXAMINATION

CANDIDATE  
NAME

CIVICS GROUP

### H2 CHEMISTRY

9729/02

Paper 2 Structured Questions

16 September 2021

2 hours

Candidates answer on the Question Paper.

Additional materials: *Data Booklet*

### READ THESE INSTRUCTIONS FIRST

Write your name and civics group in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams and graphs.

Do not use paper clips, glue or correction fluid.

The use of an approved scientific calculator is expected, where appropriate.

Answer **all** questions in the spaces provided on the Question Paper.

A Data Booklet is provided.

The number of marks is given in brackets [ ] at the end of each question or part question.

**There is one question in the paper relating to the Common Last Topic.**

**Please cancel Q5 (d) (iii) on page 19**

| Examiner's Use |     |       |
|----------------|-----|-------|
| Paper 1        | MCQ | / 30  |
| Paper 2        | Q1  | / 12  |
|                | Q2  | / 13  |
|                | Q3  | / 17  |
|                | Q4  | / 17  |
|                | Q5  | / 16  |
|                |     | / 75  |
| Paper 3        |     | / 80  |
| Paper 4        |     | / 55  |
| Total          |     | / 100 |
| Grade          |     |       |

- (a)** Describe and explain the variation in bonding of the Period 3 chlorides from  $\text{NaCl}$  to  $\text{PCl}_5$ .

.....[3]

- |                 | equation(s) | pH |
|-----------------|-------------|----|
| $\text{MgCl}_2$ |             |    |
| $\text{PCl}_5$  |             |    |

[3]

- (c) The melting point of NaCl is 801 °C and that of  $\text{MgCl}_2$  is 715 °C. Suggest an explanation for the lower than expected melting point for  $\text{MgCl}_2$ .

.....

.....

.....

.....

.....[2]

- (d) (i) Draw a dot-and-cross diagram of the  $\text{SCl}_2$  molecule. With reference to the Valence Shell Electron Pair Repulsion theory, state and explain the shape and bond angle.

.....

.....

.....

.....

.....[3]

- (ii)  $\text{OCl}_2$  molecule has a similar shape as  $\text{SCl}_2$  but a larger bond angle. Suggest why the bond angle in  $\text{OCl}_2$  is larger than that in  $\text{SCl}_2$ .

.....

.....

.....[1]

[Total: 12]



- 2 Cadmium metal and its compounds have many uses, ranging from standard cells, batteries to semiconductors. It is one of the main components used in the Weston standard cell. The simplified Weston standard cell is shown in Fig. 2.1.

The anode of this cell is cadmium while the cathode is made up of mercury with a layer of mercurous sulfate,  $\text{Hg}_2\text{SO}_4$ , deposited on top. The electrolyte is a solution of cadmium sulfate.





Fig. 2.1

At the cathode, both liquid mercury and sulfate ions are produced from mercurous sulfate. The equation for the reaction taking place at the anode is given below.



- (a) (i) State the type of reaction that takes place at the  $\text{Hg}_2\text{SO}_4/\text{Hg}$  electrode and write the half-equation for the reaction occurring at this electrode.

Type of reaction: .....

Half-equation: ..... [2]

- (ii) Hence, write the overall equation for the cell reaction.

.....[1]

- (iii) Draw on Fig. 2.1 the direction of electron flow when the reaction in (a)(ii) is occurring.

[1]

(b) The cell is capable of producing an e.m.f of 1.018 V.

- (i) Given that the  $E^\ominus$  for  $\text{Cd}^{2+}/\text{Cd}$  is  $-0.403\text{ V}$ , calculate the  $E^\ominus$  value for the cathode half-cell.

[1]

- (ii) Suggest and explain what happens to the standard cell potential  $E^\ominus_{\text{cell}}$  when

- solid  $\text{Cd}(\text{NO}_3)_2$  is added to the electrolyte.

.....

.....

.....

.....

.....

- the mass of cadmium electrode is increased.

.....

.....

.....

.....

.....

[3]



(iii) Calculate  $\Delta G^\circ$  for the Weston standard cell.

[2]

(iv) The standard enthalpy change for the same reaction in the Weston standard cell is  $-194.6 \text{ kJ mol}^{-1}$  at 298 K. Calculate  $\Delta S^\circ$  for the reaction.

[1]

(v) By considering the entropy and enthalpy changes during the reaction, comment qualitatively on the feasibility of the reaction with varying temperature.

[Calculation is **not** required for this question.]

.....  
 .....  
 .....[1]


(c) Suggest one hazard associated with the use of this standard cell.

.....  
 .....[1]

[Total: 13]



- 3 Cumene is a constituent of crude oil and coal tar. It is also used as a starting material to make other chemicals.




cumene

- (a) State the type of hybridisation of the carbon atoms in the benzene ring of cumene. Draw a labelled diagram to show the orbitals overlap that form the  $\sigma$  and  $\pi$  bonds between a pair of carbon atoms in the ring.

Type of hybridisation: .....

[2]

- (b) When cumene is reacted with bromine in the presence of ultraviolet light, a mixture of products may be obtained. This mixture includes compounds **A** and **B** with molecular formula  $C_9H_{11}Br$ .



- (i) Name and outline the mechanism of the reaction between cumene and bromine that forms compound **A**.

[4]

- (ii) Suggest what is the ratio of compound **A** to compound **B** in the mixture.

.....[1]

- (iii) In reality, compound **B** is formed in a higher proportion than compound **A**. Suggest a reason why this may be so.

.....  
 .....[1]


- (iv) State the isomeric relationship between compounds **A** and **B**.

.....[1]



(c) Compound **A** can be converted into compounds **B** and **C** as shown below in two steps.

State the reagents and conditions for each of the steps. Show the structure of the intermediate organic compound in each case.




Reagents and conditions

step 1 .....

step 2 .....

[3]





Reagents and conditions

step 1 .....

step 2 .....

[3]

(iii) Compound **D** shown below is an analogue of compound **A**.



Both compounds **A** and **D** can undergo nucleophilic substitution to form alcohols.


Deduce, with a reason, whether compound **D** would react more or less readily than compound **A**.

.....  
 .....[1]

(d) Compound **E** ( $C_9H_{12}$ ) is an isomer of cumene.

When compound **E** is treated with hot acidified  $KMnO_4$ , the organic product formed is benzene-1,3-dicarboxylic acid.

Draw the structure of compound **E**.



[1]


[Total: 17]

**Question 4 starts on the next page**



- 4 Amino acids are the building blocks of proteins which are vital for physiological functions.

Glutamic acid is an amino acid found in soy beans. The distinctive salty taste of soy sauce is due to salts of glutamic acid formed during fermentation in the production process.



There are three  $pK_a$  values associated with glutamic acid: 2.19, 4.25 and 9.67, with the  $\alpha$ -carboxylic acid having the lowest  $pK_a$  value.

- (a) State and explain the relative acidity of the two carboxylic acid functional groups in glutamic acid.

.....

.....


.....

.....


.....

.....[2]

- (b) At low pH, glutamic acid exists in its fully protonated form as shown.



When  $0.100 \text{ mol dm}^{-3}$  of glutamic acid in its fully protonated form is titrated with aqueous NaOH, the following curve is obtained.



**Fig. 4.1**

- (i) Calculate a value for the pH at point **A** in Fig. 4.1.

In this case, you may assume that the fully protonated form of glutamic acid behaves as a weak monobasic acid.

Give your answer to 1 decimal place.

[2]

- (ii) There are two predominant species present in equal amounts at point **B** in Fig. 4.1.

Draw the structures of these two species and state the pH of the solution at this point.

pH of solution = .....

[3]

- (iii) The isoelectric point of an amino acid refers to the pH at which the amino acid has no overall charge. On the graph in Fig. 4.1, indicate a point **C** which corresponds to the isoelectric point of glutamic acid.

[1]

- (c) Some properties of the amino acid alanine and 3-aminobutan-1-ol are given in Table 4.2.

**Table 4.2**

|                                                                                             | $M_r$ | solubility in water | melting point/°C |
|---------------------------------------------------------------------------------------------|-------|---------------------|------------------|
| CH <sub>3</sub> CH(NH <sub>2</sub> )CO <sub>2</sub> H<br>alanine                            | 89    | soluble             | 300              |
| CH <sub>3</sub> CH(NH <sub>2</sub> )CH <sub>2</sub> CH <sub>2</sub> OH<br>3-aminobutan-1-ol | 89    | soluble             | −1               |

- (i) Despite having a similar molecular mass, alanine and 3-aminobutan-1-ol have very different melting points. Draw the structure of alanine which accounts for the difference in the melting points.

[1]

- (ii) A  $0.100 \text{ mol dm}^{-3}$  aqueous solution of 3-aminobutan-1-ol has a pH of about 11.

Explain each of the following observations, with the aid of a relevant equation, showing the chemistry of the resultant species.

[Calculation is **not** required for this question.]

- I When the aqueous solution of 3-aminobutan-1-ol is mixed with an equal volume of  $0.100 \text{ mol dm}^{-3} \text{ HCl}$ , the pH of the resultant solution was below 7.

.....

.....

.....

.....

.....

- II When the aqueous solution of 3-aminobutan-1-ol is mixed with an equal volume of  $0.050 \text{ mol dm}^{-3} \text{ HCl}$ , the resultant solution was able to resist pH changes upon addition of a small amount of NaOH.

.....

.....

.....

.....

.....

[4]


- (iii) Give one example of a buffer system in blood.

..... [1]



- (d) Protein chains contain a large number of amino acids bonded by peptide or amide bonds. The sequence of amino acids forms the primary structure of proteins.

The hormone insulin is an example of a protein. A part of the insulin molecule is shown below.



- (i) What reagents and conditions could you use to hydrolyse the protein chain into its constituent amino acids?

.....[1]

- (ii) Draw the structural formula for **two** amino acids with the longest non-aromatic side chain that would be produced by hydrolysing this segment of the protein chain using the reagents and conditions you have given in (d)(i).


[2]

[Total: 17]



- 5 Ultraviolet-visible spectroscopy is a method to analyse concentrations of chemical compounds that can absorb a wavelength of light from the ultraviolet and visible region of the electromagnetic spectrum.

The simplified diagram below shows how the concentration of a sample can be measured using ultraviolet-visible spectroscopy.



**Fig. 5.1**

The dissolved sample is placed into the vial in which light of a single wavelength will emit from the source and pass through the sample. The distance which the light travels through the sample is known as the path length. Eventually, the detector detects the remaining intensity of light and measures the absorbance.

Beer-Lambert's Law states that for a dilute solution, the absorbance,  $A$ , of the analyte is proportional to its concentration.

$$A = \epsilon c l$$

where  $\epsilon$  is a constant measured in  $\text{M}^{-1} \text{cm}^{-1}$  ( $1 \text{ M} = 1 \text{ mol dm}^{-3}$ ),

$c$  is the concentration in  $\text{mol dm}^{-3}$ ,


$l$  is the path length in centimeter, cm.

- (a) With reference to Fig. 5.1, suggest a value of the path length in cm.

Path length: ..... cm

[1]

Buckminsterfullerene,  $C_{60}$ , is a molecule with 60 carbon atoms arranged in a spherical structure.



buckminsterfullerene,  $C_{60}$

- (b) To prepare the sample of  $C_{60}$  for analysis,  $10\text{ cm}^3$  of concentrated  $C_{60}$  was diluted to  $200\text{ cm}^3$  with a suitable solvent. The diluted  $C_{60}$  is then analysed using ultraviolet-visible spectroscopy at a specific wavelength and the absorbance value is found to be 0.96.

The value of  $\epsilon$  is  $1.75 \times 10^5\text{ M}^{-1}\text{ cm}^{-1}$ . With the aid of the Beer-Lambert's Law, calculate the concentration of the concentrated  $C_{60}$  in  $\text{mol dm}^{-3}$ .

[3]

- (c) Fig. 5.2 shows that there is deviation from Beer-Lambert's Law at high concentration.




Fig. 5.2


Suggest a reason for this observation.

.....

.....[1]

$C_{60}$  has been widely researched on its modification to give other fullerene derivatives such as *exofullerenes*, *endofullerenes* and *heterofullerenes* as shown in Fig. 5.3.

*Exofullerenes* have additional atoms or ions attached to the outer spheres while *endofullerenes* have them caged within the inner sphere. *Heterofullerenes* are formed when at least one carbon atom is replaced by another element.



**Fig. 5.3**

(d)  $C_{59}M$  is a *fullerene* derivative formed by reacting the ligand  $C_{60}$  with  $M^{n+}$ , where  $M^{n+}$  can be a metal ion such as  $Al^{3+}$  or  $Ni^{2+}$ .

(i) With reference to Fig. 5.3, state which of the three types of *fullerene* derivatives does  $C_{59}M$  belongs to.

.....[1]


(ii) State the electronic configuration of  $Ni^{2+}$ .

.....[1]

(iii) Explain why complexes formed with  $Al^{3+}$  are usually not coloured whereas those containing  $Ni^{2+}$  are coloured.

.....[1]

- (e)  $C_{60}O_3$  is an *exofullerene* which can undergo decomposition in methylbenzene to give  $C_{60}O$  and oxygen gas. The rate of decomposition of  $C_{60}O_3$  is measured using the ultraviolet-visible spectroscopy. An absorbance versus time graph is plotted as shown in Fig. 5.4.



**Fig. 5.4**

- (i) Use the graph in Fig. 5.4 to deduce the order of reaction with respect to  $C_{60}O_3$ . Hence, write a rate equation for the reaction.

.....

.....

.....[3]

- (ii) Calculate the rate constant of the reaction, stating its units.

[2]

(iii) State the effect, if any, on the half-life of  $C_{60}O_3$  when

I the concentration of  $C_{60}O_3$  is halved.

.....

II the reaction temperature is increased.

.....

[2]

(iv) Apart from absorbance, state one other variable that could be measured over time to monitor the rate of this reaction.

State how this variable could be measured in a laboratory.

.....

.....[1]

[Total: 16]



**BLANK PAGE**

