

Tutorial 6: Reaction Kinetics – Suggested Answers

- 8 (a) (I) The order of reaction with respect to a given reactant is the power to which the concentration of that reactant is raised in the rate equation.
(II) The rate constant of a reaction is the constant of proportionality in the rate equation of the reaction.
(III) The half-life of a reaction is the time taken for the concentration of a reactant to decrease to half its initial value.

- (b) Comparing experiments I and II,
when initial [sucrose] $\times (0.15/0.10 =) 1.5$, initial rate $\times (0.036/0.024 =) 1.5$
 $\Rightarrow \text{rate} \propto [\text{sucrose}]$, i.e. reaction is first order with respect to sucrose.

Comparing experiments I and III,
when initial [HCl] $\times 2$, initial rate $\times 2$
 $\Rightarrow \text{rate} \propto [\text{HCl}]$, i.e. reaction is first order with respect to HCl.

$\therefore$ rate equation: $\text{rate} = k[\text{sucrose}][\text{HCl}]$

Using data from experiment I, $k = \frac{0.024}{0.10 \times 0.10} = \underline{2.40 \text{ mol}^{-1} \text{ dm}^3 \text{ s}^{-1}}$

(data from experiment II or III can be used as well)

- (c) $\text{rate} = k[\text{sucrose}][\text{HCl}]$
 $0.048 = 2.40(y)(0.25) \quad \therefore y = 0.08$

- (d) $\text{rate} = k[\text{sucrose}][\text{HCl}]$

Since HCl is a catalyst in the reaction, its concentration is effectively constant during the reaction. The rate equation can be simplified to:

$$\text{rate} = k'[\text{sucrose}], \text{ where } k' = k[\text{HCl}].$$

The reaction becomes a pseudo first-order reaction with constant half-life:

$$t_{1/2} = \frac{\ln 2}{k'} = \frac{\ln 2}{k[\text{HCl}]} = \frac{\ln 2}{k(0.10)} = 3.0 \text{ s}$$

For experiment II, [HCl] is the same as experiment I.

Hence, $t_{1/2}$ of sucrose in experiment II

$= t_{1/2}$ of sucrose in experiment I

$= \underline{3.0 \text{ s}}$

For experiment III, [HCl] is double that of experiment I.

Hence, $t_{1/2}$ of sucrose in experiment III $= t_{1/2} = \frac{\ln 2}{k'} = \frac{\ln 2}{k(2 \times 0.10)} = \frac{3.0}{2} = \underline{1.5 \text{ s}}$

- 9 Compare experiments 1 and 2:
When $[\text{NO}_2] \times 2$, rate $\times 2$
 $\Rightarrow \text{rate} \propto [\text{NO}_2]$, i.e. reaction is first order with respect to NO_2 .

Let rate = $k[\text{NO}_2][\text{SO}_2]^m$ where m is the order of reaction with respect to SO_2 .

Compare experiments 2 and 3:

$$\frac{\text{rate}_3}{\text{rate}_2} = \frac{k(0.040)(0.40)^m}{k(0.020)(0.20)^m} = \frac{8}{2}$$

Solving, $m = 1$

So rate = $k[\text{NO}_2][\text{SO}_2]$

Alternative presentation

Compare experiments 2 and 3:

When $[\text{NO}_2] \times 2$ and $[\text{SO}_2] \times 2$, rate $\times 4$

But since reaction is first order with respect to NO_2 , it means that

rate $\times 2$ due to $[\text{NO}_2] \times 2$ AND

rate $\times 2$ due to $[\text{SO}_2] \times 2$

$\Rightarrow \text{rate} \propto [\text{SO}_2]$, i.e. reaction is first order with respect to SO_2 .

Since $[\text{SO}_2] \gg [\text{NO}_2]$ in all 3 experiments, this is a pseudo first-order reaction and rate = $k'[\text{NO}_2]$, where $k' = k[\text{SO}_2]$.

$$t_{1/2} = \frac{\ln 2}{k'} = \frac{\ln 2}{k[\text{SO}_2]}$$

Since the $[\text{SO}_2]$ in expt 1 and 2 are the same, the half-life should be the same at 48 s.
Since $[\text{SO}_2]$ in expt 3 is twice that in expt 1, the half-life would be halved to 24 s.

10 Ans: B

As the decomposition is a first order reaction, $t_{1/2} = \frac{\ln 2}{k}$.

$t_{1/2}$ is independent of $[\text{H}_2\text{O}_2]$, i.e. doubling $[\text{H}_2\text{O}_2]$ from 0.1 mol dm^{-3} to 0.2 mol dm^{-3} has no effect on $t_{1/2}$. Since $\ln 2$ and k are constants, $t_{1/2}$ is constant and likewise, time taken for H_2O_2 to decompose by 10 % will also be constant at 5 min.

- 11 (a) Since rate of reaction $\propto \frac{\Delta[\text{I}_2]}{\Delta t} \propto \frac{\text{volume of } \text{I}_2(\text{aq}) \text{ used}}{\text{time taken for solution to turn colourless}}$, by calculating $\frac{\text{volume of } \text{I}_2(\text{aq}) \text{ used}}{\text{time taken for solution to turn colourless}}$, we will be able to compare the rate of reaction for following experiments.

Expt. No.	vol. of CH_3COCH_3 / cm^3	vol. of I_2 / cm^3	vol. of $\text{H}_2\text{SO}_4(\text{aq})$ / cm^3	vol. of water / cm^3	time taken for solution to decolourise / min	$\frac{\text{vol of } \text{I}_2}{\text{time taken}} / \text{cm}^3 \text{ min}^{-1}$
1	8.0	4.0	8.0	0.0	1.0	4.0
2	8.0	4.0	4.0	4.0	2.0	2.0
3	4.0	4.0	8.0	4.0	2.0	2.0
4	8.0	2.0	8.0	2.0	0.5	4.0

Comparing experiments 1 and 2,
when volume of $\text{H}_2\text{SO}_4 \times 0.5$ (corresponding to $[\text{H}^+] \times 0.5$), rate $\times 0.5$.
 $\Rightarrow \text{rate} \propto [\text{H}^+]$, i.e. reaction is first order with respect to H^+ .

Comparing experiments 1 and 3,
When volume of $\text{CH}_3\text{COCH}_3 \times 0.5$ (corresponding to $[\text{CH}_3\text{COCH}_3] \times 0.5$, rate $\times 0.5$.
 $\Rightarrow \text{rate} \propto [\text{CH}_3\text{COCH}_3]$, i.e. reaction is first order with respect to CH_3COCH_3 .

Comparing experiments 1 and 4,
When volume of $\text{I}_2 \times 0.5$ (corresponding to $[\text{I}_2] \times 0.5$, rate remains unchanged.
 $\Rightarrow \text{rate} \propto [\text{I}_2]^0$, i.e. reaction is zero order with respect to I_2 .

- (b) (I) The rate equation for a reaction is a mathematical expression that relates the rate of reaction to the concentration of each reactant raised to the appropriate power.
 (II) The rate-determining step is the slowest step in the reaction mechanism of a multi-step reaction and it determines the overall reaction rate.

(c) $\text{rate} = k[\text{CH}_3\text{COCH}_3][\text{H}^+]$; Overall order = 2

(d) In mechanism **A**, the slow step (also the rate-determining step) is a bimolecular elementary reaction involving one molecule of CH_3COCH_3 and one H^+ ion. The rate equation is $\text{rate} = k'[\text{CH}_3\text{COCH}_3][\text{H}^+]$, which is similar to the rate equation in (c).

⇒ Mechanism A fits the observed kinetic data.

In mechanism **B**, the slow step is a bimolecular elementary reaction involving one molecule of CH_3COCH_3 and one molecule of I_2 . The rate equation is $\text{rate} = k''[\text{CH}_3\text{COCH}_3][\text{I}_2]$, which is different from the rate equation in (c).

⇒ Mechanism **B** is inconsistent with the observed kinetic data.

[Note: The reaction is zero order with respect to I_2 , so I_2 is only involved in the fast step (which is not rate-determining).]

(e) As the reaction proceeds by a similar mechanism, bromine (like iodine) is not involved in the slow/rate-determining step. The reaction is therefore zero order with respect to bromine. Hence, the rate of reaction is similar to that of the reaction between propanone and iodine under the same experimental conditions.

(Note: Even though the Br-Br bond is stronger than the I-I bond, the rate of reaction will be similar, because slow/rate-determining step does not involve the breaking of the halogen-halogen bond.)

12 (a) amount of $\text{S}_2\text{O}_3^{2-}$ used = $0.050 \times 0.0020 = 0.000100 \text{ mol}$



amount of $\text{I}_2 = \frac{1}{2} \times 0.000100 = \underline{5.00 \times 10^{-5} \text{ mol}}$

(b) $[\text{I}_2] = \frac{\text{amount of iodine}}{\text{volume of mixture}} = \frac{5.00 \times 10^{-5}}{0.100} = 5.00 \times 10^{-4} \text{ mol dm}^{-3}$

Initial rate of reaction $\approx \frac{[\text{I}_2]}{t}$

Expt No	Time for the appearance of deep blue colour/ s	Initial rate $\approx \frac{[\text{I}_2]}{t} / \text{mol dm}^{-3} \text{ s}^{-1}$
1	33	1.52×10^{-5}
2	100	5.00×10^{-6}
3	67	7.46×10^{-6}
4	50	1.00×10^{-5}

Comparing experiments 2 and 4,
 when volume of $\text{H}_2\text{O}_2 \times 2$ (corresponding to $[\text{H}_2\text{O}_2] \times 2$), initial rate $\times 2$.
 ⇒ $\text{rate} \propto [\text{H}_2\text{O}_2]$
 ⇒ reaction is first order with respect to H_2O_2 .

Comparing experiments 2 and 3,
 when volume of KI $\times 1.5$ (corresponding to $[\text{KI}] \times 1.5$), initial rate $\times 1.5$.
 ⇒ $\text{rate} \propto [\text{KI}]$
 ⇒ reaction is first order with respect to KI.

Let rate equation be $\text{rate} = k[\text{H}_2\text{O}_2][\text{KI}][\text{HCl}]^b$ where b is the order of reaction with respect to HCl .

Comparing experiments 1 and 2 and using the fact that $[\text{X}] \propto V_{\text{X}}$ since total volume is a constant,

$$\frac{\text{Rate}_1}{\text{Rate}_2} = \frac{k(15)(10)(5)^b}{k(5)(10)(10)^b} = \frac{1.52 \times 10^{-5}}{5.00 \times 10^{-6}} \Rightarrow 3(0.5)^b = 3.04 \Rightarrow b = 0$$

$\Rightarrow$ reaction is zero order with respect to HCl .

(c) $\text{rate} = k[\text{H}_2\text{O}_2][\text{KI}]$; units of rate constant is $\text{mol}^{-1} \text{dm}^3 \text{s}^{-1}$.

(d) $y = 33 \text{ s}$.

Explanation (not required by question):

The total volume is twice that of all the other experiments.


The volume of each reagent used is twice that of Experiment 1

$\Rightarrow$ initial concentrations of all reactants are the same as that of Experiment 1

$\therefore$ time taken is the same as Experiment 1.

13

(i) Use the graphs to calculate the initial rates of reaction during the two runs.



$$\text{For run A, initial rate} = -\frac{0.0010 - 0}{0 - 32} = \underline{3.125 \times 10^{-5}} = \underline{3.13 \times 10^{-5} \text{ mol dm}^{-3} \text{ min}^{-1}}$$

$$\text{For run B, initial rate} = -\frac{0.0010 - 0}{0 - 16} = \underline{6.25 \times 10^{-5} \text{ mol dm}^{-3} \text{ min}^{-1}}$$

(Note: On the graph, label coordinates used to obtain the initial rates.)

(ii) Comparing run A and run B,
when $[\text{OH}^-] \times 2$, initial rate $\times 2$
 $\Rightarrow \text{rate} \propto [\text{OH}^-]$
 $\Rightarrow$ reaction is first order with respect to OH^-

(iii) From the graph of run B,
First $t_{1/2} = \text{Second } t_{1/2} = 11 \text{ min} \Rightarrow \underline{t_{1/2} \text{ is constant}}$
 $\therefore$ reaction is first order with respect to ester.

[Note: Show all construction lines and label at least two $t_{1/2}$ on the graph (refer to the above graph in **(b)(i)**)]

[For run B, $[\text{OH}^-]$ is $\frac{0.40}{0.0010} = 400$ times that of [ester] and $[\text{OH}^-]$ remains effectively constant during the reaction. Hence the reaction follows pseudo first-order kinetics.]

(iv) $\text{rate} = k[\text{ester}][\text{OH}^-]$

(v) From run B, $6.25 \times 10^{-5} = k(0.0010)(0.40)$
 $\therefore k = \underline{0.156 \text{ mol}^{-1} \text{ dm}^3 \text{ min}^{-1}}$

OR

From run A, $3.125 \times 10^{-5} = k(0.0010)(0.20)$
 $\therefore k = \underline{0.156 \text{ mol}^{-1} \text{ dm}^3 \text{ min}^{-1}}$

(vi) Since the reaction follows pseudo first-order kinetics.

$\text{rate} = k'[\text{ester}]$, where $k' = k[\text{OH}^-]$ and $t_{1/2} = 11 \text{ min}$ (from **(iii)**)

$$t_{1/2} = \frac{\ln 2}{k'} \Rightarrow k' = \frac{\ln 2}{t_{1/2}} \Rightarrow k[\text{OH}^-] = \frac{\ln 2}{t_{1/2}}$$

$$\therefore k = \frac{\ln 2}{t_{1/2}[\text{OH}^-]} = \frac{\ln 2}{(11)(0.40)} = \underline{0.158 \text{ mol}^{-1} \text{ dm}^3 \text{ min}^{-1}}$$

- 14 (a) (i) [B] remains approximately constant as the reaction proceeds. The rate of reaction is only dependent on [A] and the order of reaction with respect to A can thus be found.

Comments:

- Many students realised that A was the limiting reagent when B was added in excess. However, the emphasis in this question is on "*large excess*". Examiners are looking for an answer that is related to how we study the kinetics of the reaction.
- Examiners accepted answers that described a pseudo first-order reaction. However, in actual fact, when B is added in excess, we do not yet know if the reaction is a pseudo first-order or a pseudo second-order reaction. It would be more accurate to simply describe the reaction as a pseudo order reaction.

- (ii) If the reaction were to go to completion, final $[\text{C}] = \text{initial } [\text{A}]$
 $= \underline{0.12 \text{ mol dm}^{-3}}$

First $t_{1/2}$ is time taken for $[\text{C}]$ to increase from 0 to 0.06 mol dm^{-3} while second $t_{1/2}$ is time taken for $[\text{C}]$ to increase from 0.06 to 0.09 mol dm^{-3} .


From the graph, 1st $t_{1/2} = 3.7 \text{ min}$ and 2nd $t_{1/2} = 3.7 \text{ min}$
 Since 1st $t_{1/2} = 2^{\text{nd}} t_{1/2}$, the reaction is first order with respect to A.

Note:

- The final $[\text{C}]$ should be used to determine $t_{1/2}$.
- Show all construction lines and label at least two $t_{1/2}$ on the graph (refer to graph in solutions to **(b)(i)** on the next page)

Comments:

- The final [C] should be calculated using the stoichiometric ratios of the equation given and the calculation should take into account the fact that A is the limiting reagent. In this case, the final [C] cannot be obtained from the graph.
- Students should learn how to find the half-life of a reaction using a [product] vs time graph. The half-lives should also be clearly marked out on the graph, and the values written in the space provided for the answer. Quite a number of students stated the 2nd half-life is 7.4 min, which is wrong. 7.4 min is the sum of the first two half-lives.

**Comments:**

- Generally well done, but a few students forgot to include the point at the origin.
- Use a sharp pencil to plot the points accurately.

(ii) **Method 1:** Compare initial rates of experiments 1 and 2.

$$\text{initial rate of experiment 1} = g_1 = \frac{0.090 - 0}{4.0 - 0} = 0.0225 \text{ mol dm}^{-3} \text{ min}^{-1}$$

$$\text{initial rate of experiment 2} = g_2 = \frac{0.045 - 0}{4.0 - 0} = 0.01125 \text{ mol dm}^{-3} \text{ min}^{-1}$$

Since initial rate is halved when [B] is halved, the reaction is first-order with respect to B.

(Note: On the graph, label coordinates used to obtain g₁ and g₂)

Method 2: Compare half-lives of experiments 1 and 2.

Let order of reaction with respect to B be y.

$$\text{Rate} = k[A][B]^y = k'[A] \text{ where } k' = k[B]^y$$

Since B is used in large excess, $[B]^y$ is approximately constant such that the reaction is a pseudo first-order reaction.

$$t_{1/2} = \frac{\ln 2}{k'} = \frac{\ln 2}{k[B]^y} \Rightarrow t_{1/2} \propto \frac{1}{[B]^y} \Rightarrow t_{1/2} \times [B]^y \text{ is a constant}$$

For experiment 1, $t_{1/2} = 3.7 \text{ min}$ and $[B] = 2.00 \text{ mol dm}^{-3}$

For experiment 2, $t_{1/2} = 7.4 \text{ min}$ and $[B] = 1.00 \text{ mol dm}^{-3}$

$$3.7(2.00)^y = 7.4(1.00)^y$$

$$\frac{3.7}{7.4} = \left(\frac{1.00}{2.00} \right)^y \Rightarrow y = 1 \Rightarrow \text{reaction is first order with respect to B}.$$

Comments:

- The initial rate is obtained by drawing a tangent to a curve at $t = 0$ and finding the gradient of that tangent. Some students chose to draw a tangent at some value of t for which $t > 0$ and found the instantaneous rate instead. Students who used instantaneous rate were unable to arrive at the correct answer as the calculations were more difficult.
- Students should show, on the graph, the construction lines for the tangent. Students also need to show the examiners how the value of the gradient was calculated. Note that the unit for time is in minutes (min).
- Students should be mindful of how they use words such as “increased by 2 times”. When 3 increases by 2 times, it becomes $3 + 2(3) = 9$, not 6. When 3 is doubled, the result is 6.
- Students who used **Method 2** often did not show clearly how the half-life is related to $[B]$.
- Obtaining a constant half-life for the reaction in experiment 2 meant that the reaction was first-order with respect to A, not first-order with respect to B.

(iii) $\text{rate} = k[A][B]$

Comments:

- Some students gave wrong answers such as “rate equation = $k[A][B]$ ” and “ $k = [A][B]$ ”.

(iv) **Method 1:** Use initial rate

Using results from experiment 1 in (b)(ii),

initial rate of expt 1 = $0.0225 \text{ mol dm}^{-3}$

Since $\text{rate} = k[A][B]$, at $t = 0 \text{ min}$, $0.0225 = k(0.12)(2.00)$

$$\Rightarrow k = \frac{0.0225}{(0.12)(2.00)} = \underline{0.0938 \text{ mol}^{-1} \text{ dm}^3 \text{ min}^{-1}}$$

Method 2: half-life

Using results from experiment 1 in (b)(ii),

$$t_{1/2} = \frac{\ln 2}{k'} = \frac{\ln 2}{k(2.00)} = 3.7$$


$$k = \frac{\ln 2}{(3.7)(2.00)} = \underline{0.0937 \text{ mol}^{-1} \text{ dm}^3 \text{ min}^{-1}}$$

Comments:

- Many students did not realise that time was in minutes rather than seconds.
- Students who prefer using the half-life method should bear in mind that that $k' = k[B]$, as the reaction given is a pseudo first-order reaction. The required rate constant is k , not k' .

- (c) (i) A catalyst increases the rate of a reaction by providing an alternative reaction pathway of lower activation energy.

In the catalysed reaction, since the activation energy (E_a (catalysed reaction)) is lower than that of the uncatalysed reaction (E_a), more reactant particles have energy greater than or equal to the activation energy than in the uncatalysed reaction. This is shown by the larger shaded area for the catalysed reaction in the diagram below.




The frequency of effective collision increases accordingly, and hence the reaction rate increases due to a larger rate constant.

Comments:

- This is a simple “recall question”. Students must learn how to sketch the Maxwell-Boltzmann distribution curves properly and use them to explain how the rate of a reaction is affected by either a change in temperature or the use of a catalyst.
- In this case, there is only one curve. The asymmetric curve must start from the origin and the asymptotic behaviour must be shown. Indicate the legend clearly.

- (ii) As temperature increases from T_1 to T_2 K, the average kinetic energy of the reactant particles increases. As such, significantly more reactant particles have energy greater than or equal to the activation energy of the reaction. This is shown by the significantly larger shaded area at a higher temperature in the diagram below.




Consequently, the frequency of effective collisions increases accordingly, and hence the reaction rate increases due to a larger rate constant.

- 15** The reaction is an autocatalytic reaction with the H^+ ions, produced from the dissociation of CH_3CO_2H , acting as the autocatalyst.

Initially, the reaction is slow as it has a high activation energy.


The reaction rate then increases as the H^+ ions produced act as the autocatalyst, providing an alternative pathway with lower activation energy.

Towards the end of the reaction, the concentrations of the reactants are very low, so the reaction rate decreases even though there is an adequate supply of catalyst.



- 16** (a) The substrate and the enzyme molecules have complementary shapes so that they fit together like a key into its lock. The specific shape of the enzyme gives the enzyme its specificity as only substrate of a specific shape can interact with it. When enzymes and substrate molecules collide in the correct orientation, the substrate (reactant) fits the active site on the enzyme and is bound by attractive forces at the active site. An enzyme-substrate complex is formed, and this weakens the bonding within the substrate. Once the catalysed reaction has occurred, the enzyme-substrate complex breaks apart as the product formed is no longer of the right shape to fit into the active site.

(b) (i)



- (ii) Reaction rate increases with temperature only until the optimal temperature of the enzyme is reached.

Beyond the optimal temperature, this can result in a change in the three-dimensional conformation of the active site which then causes the enzyme to become denatured. The substrate is no longer complementary to the conformation/shape of active site of the enzyme, lowering the rate of reaction.

(c) (i) pH 3

(ii) At pH 8, the shape of the active site changes and is no longer complementary in shape to the substrate, hence it cannot bind to the substrate and no catalysis takes place.